

CANINE VACCINATION PROTOCOLS

DHPP

1. Begin at 6 weeks or earliest possible thereafter
2. Minimum of 2 vaccinations
3. Repeat at 3 week intervals to a minimum of 16 weeks of age
4. Booster dose 1 year later.
5. Booster doses thereafter every 3 years (usually at 4, 7, 10 years of age)

Rabies

1. All canines at 3 months or soonest possible afterwards
2. May not administer within 10 days of a human bite incident
3. One year later, booster
4. After that, booster dose every 3rd year
5. If exposure (fight) with vector animal (bat, skunk, fox, raccoon) get booster dose

Lyme

1. At risk canines at 12 weeks of age or as soon after as possible
2. Booster 2-3 weeks later for anamnestic response
3. Booster dose yearly

Bordetella

1. At risk canines at 8 weeks of age
2. Booster dose 2-3 weeks later
3. Booster dose yearly

Parainfluenza

1. At risk canines at 8 (minimum) weeks of age
2. Booster dose 2-4 weeks later
3. Booster dose 1 year later
4. Booster dose every 3 years

Leptospirosis Bacterin

1. At risk canines at 12 weeks of age
2. Booster dose 2-3 weeks later
3. Booster dose yearly

Porphyromonas Bacterin

1. At risk canines when adult teeth are in (or at veterinarian's recommendation)
2. Booster dose 2-3 weeks later
3. Booster dose at 6 or 12 month intervals, depending upon veterinarian's recommendation
4. Not recommended to administer with other vaccinations in dogs weighing less than 20 pounds as the pet is likely to have muscle aches, fever afterwards.